

A novel approach to quantifying the sensitivity of current and future cosmological datasets to the neutrino mass ordering through Bayesian hierarchical modeling

Martina Gerbino,^{1,2,*} Massimiliano Lattanzi,^{3,†} Olga Mena,⁴ and Katherine Freese^{1,2,5}

¹*The Oskar Klein Centre for Cosmoparticle Physics, Department of Physics, Stockholm University, AlbaNova, SE-106 91 Stockholm, Sweden*

²*The Nordic Institute for Theoretical Physics (NORDITA), Roslagstullsbacken 23, SE-106 91 Stockholm, Sweden*

³*Dipartimento di Fisica e Scienze della Terra, Università di Ferrara and INFN, Sezione di Ferrara, Polo Scientifico e Tecnologico - Edificio C Via Saragat, 1, I-44122 Ferrara, Italy*

⁴*IFIC, Universidad de Valencia-CSIC, 46071, Valencia, Spain*

⁵*Michigan Center for Theoretical Physics, Department of Physics, University of Michigan, Ann Arbor, MI 48109, USA*

(Dated: November 24, 2016)

We present a novel approach to derive constraints on neutrino masses, as well as on other cosmological parameters, from cosmological data, while taking into account our ignorance of the neutrino mass ordering. We derive constraints from a combination of current as well as future cosmological datasets on the total neutrino mass M_ν and on the mass fractions $f_{\nu,i} = m_i/M_\nu$ (where the index $i = 1, 2, 3$ indicates the three mass eigenstates) carried by each of the mass eigenstates m_i , after marginalizing over the (unknown) neutrino mass ordering, either normal ordering (NH) or inverted ordering (IH). The bounds on all the cosmological parameters, including those on the total neutrino mass, take therefore into account the uncertainty related to our ignorance of the mass hierarchy that is actually realized in nature. This novel approach is carried out in the framework of Bayesian analysis of a typical hierarchical problem, where the distribution of the parameters of the model depends on further parameters, the *hyperparameters*. In this context, the choice of the neutrino mass ordering is modeled via the discrete *hyperparameter* h_{type} , which we introduce in the usual Markov chain analysis. The preference from cosmological data for either the NH or the IH scenarios is then simply encoded in the posterior distribution of the *hyperparameter* itself. Current cosmic microwave background (CMB) measurements assign equal odds to the two hierarchies, and are thus unable to distinguish between them. However, after the addition of baryon acoustic oscillation (BAO) measurements, a weak preference for the normal hierarchical scenario appears, with odds of 4:3 from Planck temperature and large-scale polarization in combination with BAO (3:2 if small-scale polarization is also included). Concerning next-generation cosmological experiments, forecasts suggest that the combination of upcoming CMB (CORe) and BAO surveys (DESI) may determine the neutrino mass hierarchy at a high statistical significance if the mass is very close to the minimal value allowed by oscillation experiments, as for NH and a fiducial value of $M_\nu = 0.06$ eV there is a 9:1 preference of normal versus inverted hierarchy. On the contrary, if the sum of the masses is of the order of 0.1 eV or larger, even future cosmological observations will be inconclusive. The innovative statistical strategy exploited here represents a very simple, efficient and robust tool to study the sensitivity of present and future cosmological data to the neutrino mass hierarchy, and a sound competitor to the standard Bayesian model comparison. The unbiased limit on M_ν we obtain is crucial for ongoing and planned neutrinoless double beta decay searches.

I. INTRODUCTION

According to the standard theory of neutrino oscillations (see e.g. [1] for an updated review and relevant references), the observed neutrino flavours ν_α are a superposition of the massive eigenstates ν_i :

$$|\nu_\alpha\rangle = \sum_i U_{\alpha i}^* |\nu_i\rangle \quad (1)$$

where the index α can be any of the three active neutrino flavours e, μ, τ , the index $i = 1, 2, 3$ runs over the

three massive eigenstates and U is the Pontecorvo-Maki-Nakagawa-Sakata mixing matrix, containing the neutrino mixing angles as well as the CP violating phases (one Dirac phase, as well as two additional Majorana phases, that are non vanishing only if neutrinos are Majorana particles).

Cosmological measurements of the cosmic microwave background (hereafter CMB) anisotropies and of the spatial distribution of galaxies provide the tightest bounds on the total neutrino mass, defined as the sum of the three neutrino mass eigenstates, i.e. $M_\nu \equiv \sum m_{\nu,i} \equiv m_1 + m_2 + m_3$. The most reliable bound that can be obtained combining Planck data with external datasets is $M_\nu < 0.21$ eV (at 95% CL) [2], from temperature plus large-scale polarization CMB anisotropies and baryon acoustic oscillation (BAO) data (see also Refs [3–7] for constraints obtained by the combination of additional

* martina.gerbino@fysik.su.se

† lattanzi@fe.infn.it

datasets and/or in more extended cosmological scenarios). From neutrino oscillation data, we know that at least two out of the three mass eigenstates should be massive, as two different mass splittings are measured with percent accuracy by current experiments: the solar $\Delta m_{21}^2 \equiv m_2^2 - m_1^2 \simeq 7.6 \times 10^{-5} \text{ eV}^2$ and the atmospheric $|\Delta m_{31}^2| \equiv |m_3^2 - m_1^2| \simeq 2.5 \times 10^{-3} \text{ eV}^2$ mass gaps. Matter effects in the sun tell us that the mass eigenstate with the larger electron neutrino fraction has the smaller mass. We identify this state with “1” and the heavier state (with a smaller electron neutrino fraction) with “2”. Therefore, the solar mass splitting is positive. However, current neutrino oscillation data are unable to determine the sign of the largest mass splitting, the atmospheric mass gap. Two possible scenarios therefore appear, corresponding to the two possible signs of Δm_{31}^2 : the normal hierarchy (NH hereafter), in which the atmospheric gap is positive, and corresponds to $m_1 < m_2 < m_3$, and the inverted hierarchy (IH in what follows), in which the atmospheric gap is negative, and corresponds to $m_3 < m_1 < m_2$ ¹. Assuming that the mass of the lightest mass eigenstate is zero, which equals to set to zero the mass of m_1 (m_3) in the NH (IH), it is possible to obtain a lower bound on the sum of neutrino masses of $M_\nu = \sqrt{\Delta m_{21}^2} + \sqrt{\Delta m_{31}^2} \simeq 0.06 \text{ eV}$ ($M_\nu = \sqrt{\Delta m_{31}^2} + \sqrt{\Delta m_{31}^2} \simeq 0.1 \text{ eV}$) from neutrino oscillation measurements.

Neutrino mixing phenomena are sensitive to the neutrino mass splittings only, not to the individual neutrino masses nor to the overall mass scale. Cosmology provides one of the most suitable places where to test and extract the neutrino mass ordering [8–11], see also the recent work of Refs. [7, 12]. Despite the fact that current bounds on the neutrino mass M_ν show a dependence on how the mass is distributed among the three mass eigenstates [6], present cosmological measurements are not able to firmly single out nature’s choice for the mass hierarchy. Consequently, in the absence of a robust measurement of the neutrino mass ordering, a desirable bound on M_ν would be one which does not rely on any assumption (or, to be more precise: that relies on the less informative possible assumption) about the hierarchical distribution of the total mass among the three eigenstates. This kind of problem, where the distribution of the parameters of the model under scrutiny are themselves conditionally dependent on the so-called *hyperparameters* (namely, the bounds on M_ν are extracted by assuming a specific mass splitting), is a typical example of a hierarchical model in statistical inference. In this work, we propose a novel method to get a hierarchy-independent bound on M_ν , by

means of a new discrete parameter, the hyperparameter, h_{type} (that can in practice be identified with the sign of the atmospheric mass splitting), introduced in the standard Monte Carlo Markov chain (MCMC) analysis. This innovative strategy benefits from the fact that the sensitivity to the neutrino hierarchy is simply and unbiasedly extracted from the posterior probability distribution of h_{type} . We shall add this parameter while analyzing current and future cosmological data, to illustrate the power of this technique.

We stress that our approach is different from the one, already found in the literature, in which a single, continuous parameter is used to parametrize both the sum of neutrino masses and the hierarchy [11]. The drawback of the approach proposed in Ref. [11], when used in an MCMC framework², is that it is not possible to disentangle the prior assumptions on the individual masses and on the hierarchy; in particular, a flat (uninformative) prior on the hierarchy implies a non-flat prior on the mass. In our approach, we are free to specify noninformative priors for both the hierarchy and the mass of the lightest eigenstate.

Apart from cosmological probes, there also exist laboratory avenues which are sensitive to the absolute mass scale. In this context, neutrinoless double β decay ($0\nu 2\beta$) searches (see e.g. Refs. [13–16]) are intriguing, as a positive signal would guarantee that neutrinos have a non-zero Majorana mass [17]. Double beta decay is a rare spontaneous nuclear transition in which the charge of two isobaric nuclei changes by two units, emitting two electrons. The dominant mode of this decay also produces two electron antineutrinos, conserving lepton number and therefore, it is allowed in the standard model framework. Double β decay without antineutrino emission, violating lepton number by two units, is the neutrinoless double β decay. Planned $0\nu 2\beta$ experiments might have the required sensitivity to completely cover the region of the parameter space where a positive signal is expected in the case of IH distribution of the total neutrino mass. Robust limits on the total neutrino mass coming from cosmology can further reduce the allowed region of the parameter space where to look for $0\nu 2\beta$ events.

The paper is organized as follows: we describe our method and provide details of the parameterization we adopt in Sec. II; we present and discuss the implications for present cosmological data, future CMB and BAO missions and neutrinoless double beta decay experiments in Secs. III, IV and V respectively. We conclude in Sec. VI.

¹ Recent results for the NOvA long baseline experiment show that the best fit is obtained for the NH scheme, and rule out at $\sim 3\sigma$ a small region of the IH scenario, for some particular ranges of the mixing parameters. However, a large fraction of the IH region is still allowed. Antineutrino data can shed light on these results, see <http://nusoft.fnal.gov/nova/results/index.html> [19].

² While this paper was being finalized, Ref. [12] appeared, using the parameterization proposed in Ref. [11] in an MCMC framework to derive constraints on the neutrino hierarchy.

II. METHOD AND DATA

A. Statistical framework and choice of the relevant parameters

The problem we deal with in this work is a typical example of a statistically hierarchical³ model (see e.g. [18]). A key feature of hierarchical problems is that the parameters $\vec{\theta}$ of the model introduced for constraining the observables through the data \vec{d} are modeled conditionally on further parameters, the *hyperparameters* $\vec{\phi}$, which have themselves their own prior probability distribution $p(\vec{\phi})$. As a result, we can define a joint prior distribution

$$\Pi \equiv p(\vec{\phi}, \vec{\theta}) = p(\vec{\phi})p(\vec{\theta}|\vec{\phi}) \quad (2)$$

so that the proper posterior distribution $\mathcal{P} \equiv p(\vec{\theta}, \vec{\phi}|\vec{d})$ of the parameters (both “normal” and “hyper”) can be written, using Bayes’ theorem, as

$$\mathcal{P} = \frac{\Pi \cdot \mathcal{L}}{\mathcal{E}} \propto p(\vec{\phi}, \vec{\theta})p(\vec{d}|\vec{\phi}, \vec{\theta}) = p(\vec{\phi}, \vec{\theta})p(\vec{d}|\vec{\theta}) \quad (3)$$

where $\mathcal{L} \equiv p(\vec{d}|\vec{\theta}, \vec{\phi})$ is the likelihood function, in which we dropped the explicit dependence on $\vec{\phi}$, since the data depends on $\vec{\phi}$ only through $\vec{\theta}$, and $\mathcal{E} \equiv \int \mathcal{L} \cdot \Pi d\vec{\theta}d\vec{\phi}$ is the model evidence, or marginal likelihood. The latter does not depend on the parameters, and thus represents just a multiplicative constant as long as parameter estimation is concerned⁴.

In the case under investigation in this work, the model parameters are extracted conditionally on the choice of the neutrino mass hierarchy. This choice is modelled by introducing a discrete hyperparameter h_{type} that can take two values, corresponding to NH and IH (i.e. to $\text{sgn}(\Delta m_{31}^2) = +1$ or -1 , respectively). Since little is known from current experiments about the preference for one of the two neutrino hierarchies, either normal or inverted, we assign equal *a priori* probability to the two possible outcomes that h_{type} could take.

We therefore perform a MCMC analysis of an eight-dimensional parameter space. We consider the usual set of six cosmological parameters in the Λ CDM scenario, namely the baryon density $\Omega_b h^2$, the cold dark matter density $\Omega_c h^2$, the angular size of the sound horizon θ_s , the reionization optical depth τ , the scalar spectral index n_s and the amplitude $\ln[10^{10} A_s]$ of the power spectrum of primordial scalar perturbations normalized at

the pivot scale $k_0 = 0.05 \text{ Mpc}^{-1}$. All these parameters are extracted from flat prior distributions.

The inclusion of massive neutrinos is performed in the following way: we assume three massive non-degenerate eigenstates sharing the same temperature $T_\nu = (4/11)^{1/3} T_\gamma$. We sample, again with a flat prior⁵, over the lightest eigenstate mass m_{light} (thus corresponding to the seventh parameter of the model), which is equivalent to m_1 (m_3) in the NH (IH) scenario. The mass of the remaining two neutrino states is set by oscillation measurements through the solar and atmospheric mass gaps, i.e. the so-called squared mass differences, defined as $\Delta m_{ij}^2 = m_i^2 - m_j^2$, with $i, j = 1, 2, 3$. While the solar mass splitting is by convention positive, i.e. $\Delta m_{21}^2 > 0$, the sign of the atmospheric mass gap Δm_{31}^2 , as previously stated, remains still unknown, and it depends on the hierarchical distribution of the total mass among the eigenstates, with $\Delta m_{31}^2 > 0 (< 0)$ in the NH (IH) scenario. This is the reason why the lightest eigenstate corresponds to m_1 in the NH scenario, while it is mapped onto m_3 in the IH scenario. We use the latest best-fit values for the oscillation mass gaps [20, 21].

As anticipated at the beginning of this section, the choice of the hierarchy is addressed via the discrete hyperparameter h_{type} , which is the eight parameter of the model. At each step, we extract h_{type} from $\{\text{NH}, \text{IH}\}$, assigning equal *a priori* probability to the two hierarchical scenarios (i.e., we use the discrete equivalent of a flat prior). In the formalism sketched at the beginning of the section, $\vec{\theta} \equiv (\Omega_b h^2, \Omega_c h^2, \theta_s, \tau, n_s, \ln[10^{10} A_s], m_{\text{light}})$, while $\vec{\phi} = h_{\text{type}}$. The inclusion of the hyperparameter allows us to handle our ignorance about the true hierarchical distribution of the mass as a nuisance parameter, to be marginalized over. In this way, the posterior distribution of h_{type} for a given datasets contains information about the preference for one of the two hierarchies from that dataset. This is easily done from the chains generated by the MCMC algorithm, by computing the marginalized probabilities \mathcal{P}_{NH} and \mathcal{P}_{IH} , defined as

$$\mathcal{P}_{\text{NH}} \equiv p(h_{\text{type}} = \text{NH}|\vec{d}) = \int \mathcal{P}(\vec{\theta}, h_{\text{type}} = \text{NH}) d\vec{\theta}, \quad (4)$$

and similarly for \mathcal{P}_{IH} . This information is conveyed by reporting the “odds” for NH vs. IH, i.e. the ratio $\mathcal{P}_{\text{NH}} : \mathcal{P}_{\text{IH}}$.

To compute the cosmological constraints and the posterior probability distributions in this extended Λ CDM scenario, we make use of the latest version of the publicly available MCMC package `cosmomc` [22, 23], exploiting the Gelman and Rubin statistics for monitoring the convergence of the generated chains [24]. We quote our results in terms of 95% credible intervals for the param-

³ The meaning of “hierarchical” here has not to be confused with the two different neutrino mass distributions, or hierarchies. While we shall make use of the same terminology to refer to different concepts throughout the text, the context in which it is employed will help solving the ambiguity.

⁴ We note that, even though in this paper we are in principle also addressing a problem of model selection - i.e., determining the correct model for neutrino hierarchy -, the use of the hyperparameter allows to map this into a parameter estimation problem.

⁵ Since the relation between m_{light} and M_ν is nonlinear, this is in principle different than sampling over M_ν , as it is usually done.

eters. Given that we will be dealing with possibly multimodal distributions, the credible intervals can consist of the union of disjointed regions.

B. Cosmological Datasets

Current CMB and BAO measurements are considered. Results are presented separately for CMB temperature and low-multipole polarization data (the former ranging from multipoles $\ell = 2$ up to $\ell = 29$) from the Planck mission (TT + lowP), and for the addition, to the previous measurements, of the high multipole (i.e. small-scale) polarization and cross-correlation spectra, i.e. the full Planck data release (TT, TE, EE + lowP), see Refs. [2, 25, 26]. We remind the reader that since, as discussed in Refs. [2, 26], small-scale polarization data could still be affected by low-level residual systematics, results obtained without using them should be regarded as more reliable. These measurements are analyzed by means of the publicly available Planck likelihood code [26], and foregrounds or extra nuisance parameters are carefully treated following the prescription detailed in Refs. [2, 26].

We combine the two CMB datasets described above with geometrical information from galaxy clustering, i.e. via the BAO signature. All the BAO measurements we exploit here are expressed as determinations of $D_V(z_{\text{eff}})/r_s(z_{\text{drag}})$, with

$$D_V(z) = \left[(1+z)^2 D_A(z)^2 \frac{z}{H(z)} \right]^{1/3} \quad (5)$$

representing a combination of the line-of-sight clustering information (as encoded by the Hubble parameter H) and the transverse clustering information (encoded in the angular diameter distance D_A) at the effective redshift z_{eff} of the survey, and $r_s(z_{\text{drag}})$ being the sound horizon at the drag epoch⁶. Concretely, we make use of the BAO results from the 6dF Galaxy Survey (6dFGS) [28] and from the BOSS DR11 LOWZ and CMASS samples [29], focusing at $z_{\text{eff}} = 0.106$, and $z_{\text{eff}} = 0.32, 0.57$, respectively. The combination of these measurements will be referred to as BAO. The addition of galaxy clustering measurements, apart from breaking pure *geometrical* degeneracies among different cosmological parameters (as, for instance, the one existing between the Hubble constant H_0 and the neutrino mass M_ν [30]), also helps enormously in pinning down the neutrino mass limits, as the free streaming nature of sub-eV neutrinos will leave a clear imprint in the galaxy power spectrum at scales in the regime of interest, see e.g. [6].

We also perform forecasts for future CMB and galaxy clustering data. For the fiducial values of the six Λ CDM

parameters, we use the mean values of the estimates reported in [31] for PlanckTT+SIMlow. Concerning CMB measurements, we consider a future CORe-like [32] satellite mission, generating mock lensed temperature and polarization data accordingly to Refs. [33, 34]. We assume perfect foreground subtraction as well as precise control of systematics. The expected noise spectra (which relies on specific experimental setup, such as the sky fraction, the beam width, and the temperature and polarization sensitivities) has been verified against previous results in the literature [35, 36], finding an excellent agreement. Future galaxy clustering data are added by means of the expected independent observations of the BAO signal along and across the line of sight from the Dark Energy Instrument (DESI) Experiment [37]. DESI observations will provide separate measurements of $H(z)r_s$ and $D_A(z)/r_s$ at a number of redshifts. This survey is expected to cover 14000 squared degrees of the sky in the redshift range $0.15 < z < 1.85$. We follow the DESI Conceptual Design Report, and also Ref. [38] for the percentual errors on both $H(z)r_s(z)$ and $D_A(z)/r_s(z)$ expected from the three types of DESI tracers, (namely, Emission Line Galaxies, Luminous Red Galaxies and High Redshift Quasars) and assume an identical 0.4 correlation coefficient between the percentual errors on $H(z)r_s(z)$ and $D_A(z)/r_s(z)$ (see e.g. [38, 39]).

Furthermore, we include, in the future cosmological data, a 1%-measurement of the Hubble constant H_0 . This is implemented in the form of a gaussian prior on H_0 in the analysis with future cosmological data. However, given the fact that the CORe sensitivity is such that it will expectedly allow to determine the value of H_0 below that precision, such a prior on the Hubble constant does not play any crucial role.

III. RESULTS FROM PRESENT COSMOLOGICAL MEASUREMENTS

In this section, we present the bounds on the neutrino mass parameters derived from different combinations of current cosmological probes, as well as discuss the sensitivity of the very same probes to the neutrino mass ordering. Table I shows the 95% CL constraints on the total neutrino mass, on the lightest neutrino mass, and on the individual neutrino masses associated to each neutrino mass eigenstate after marginalization over the h_{type} parameter. For each parameter that appears in the table, with the exception of h_{type} , we quote our results in the following way: if i) the 95% confidence interval includes one of the edges of the prior range for that parameter (this is the case for M_ν and for the individual masses), or ii) the posterior probability distribution is bimodal (this is sometimes the case for M_ν , see below), then we report the 95% confidence interval in the form [min, max]; iii) otherwise, we report the 95% confidence interval in the form (mean \pm uncertainty). In the last line of the table, we report the results for h_{type} , in the form (odds for

⁶ The drag epoch is defined as the time at which baryons are released from the Compton drag of the photons, see Ref. [27].

	TT + low P	TT, TE, EE + low P	TT + lowP + BAO	TT, TE, EE + low P + BAO
M_ν [eV]	[0.058 – 0.740]	[0.058 – 0.558]	[0.058 – 0.232]	[0.058 – 0.200]
m_{light} [eV]	< 0.244	< 0.183	< 0.0695	< 0.0577
m_1 [eV]	< 0.246	< 0.186	< 0.079	< 0.068
m_2 [eV]	[0.009 – 0.246]	[0.009 – 0.186]	[0.009 – 0.079]	[0.009 – 0.069]
m_3 [eV]	< 0.243	< 0.185	< 0.082	< 0.072
$\Omega_c h^2$	$0.1205^{+0.0046}_{-0.0045}$	0.1203 ± 0.0030	$0.1184^{+0.0026}_{-0.0027}$	0.1189 ± 0.0022
H_0 [km s ⁻¹ Mpc ⁻¹]	$65.1^{+4.0}_{-5.3}$	$65.6^{+2.8}_{-3.7}$	67.4 ± 1.2	67.2 ± 1.1
Ω_m	$0.346^{+0.081}_{-0.059}$	$0.337^{+0.052}_{-0.039}$	$0.313^{+0.016}_{-0.015}$	0.316 ± 0.014
h_{type} odds (NH:IH)	1 : 1	9 : 8	4 : 3	3 : 2

TABLE I. 95% credible intervals for the total neutrino mass, the mass of the lightest neutrino eigenstate and the individual neutrino masses, as well as for other cosmological parameters, for different combinations of current CMB and BAO data. All the bounds reported here also take into account information from oscillation measurements. In the last row, we quote the odds for the NH vs. the IH scenario.

NH : odds IH). The odds shown in the second column of Table I show how CMB temperature data alone are not sensitive to the different mass parameterizations. This is due to the broad bound that the CMB temperature data set on the total neutrino mass, $M_\nu < 0.740$ eV at 95% CL. In fact, the potential for cosmological observations to discriminate between the two mass orderings mainly relies on the capability to push the upper limit on M_ν close or even below 0.1 eV, the minimal value of the mass allowed by oscillation data in the case of IH. This is mainly due to the fact that the region $M_\nu < 0.1$ eV is only allowed in the case of normal hierarchy, so that tighter upper bounds on M_ν end up favouring the NH scenario simply because of the larger volume in parameter space available to the model. Moreover, the region of masses with $M_\nu \simeq 0.1$ eV is the one where the mass patterns predicted by NH and IH, and the resulting cosmological perturbations, differ the most. The differences are however small, given the sensitivity of present, and possibly also future, experiments, and the dominant contribution to the constraining power still comes from the sheer amount of volume in parameter space available to the two models. On the opposite, when $M_\nu \gg 0.1$ eV (i.e., $M_\nu \gg \Delta m_{21}^2, \Delta m_{31}^2$), we are in a situation in which both hierarchies effectively coincide with the degenerate scenario $m_1 \simeq m_2 \simeq m_3$, and the differences in the evolution of perturbations induced by the mass ordering are too small to have any observable consequence, given current sensitivities. Given that we find a bound from CMB temperature anisotropies and large-scale polarization $M_\nu < 0.740$ eV, most of the parameter space available, given this data, is in the “effectively degenerate” re-

gion, where the two hierarchies cannot be distinguished. Indeed, the posterior distributions for the mass fractions $f_{\nu,i} \equiv m_i/M_\nu$ for the case of CMB temperature data are clearly peaking on $f_{\nu,i} = 1/3$, as it would be in the fully degenerate scenario. Notice that the addition of small-scale CMB polarization measurements slightly improves the neutrino mass bound ($M_\nu < 0.558$ eV at 95% CL) but it does not change significantly the overall picture.

We note that the bounds we find seems to be larger, when a direct comparison is possible, than those found in Ref. [2] without the marginalization over the two possible mass orderings: compare, e.g., our bound $M_\nu < 0.740$ eV with $M_\nu < 0.715$ eV, the 95% upper bound from Planck TT+lowP in the context of the Λ CDM+ M_ν model, assuming three massive degenerate neutrinos [2]. The reason is the following. In the present analysis, a non-vanishing lower bound on the total neutrino mass, $M_{\nu,\text{min}} = 0.058$ eV is naturally imposed by taking into account neutrino oscillation measurements, while in Ref. [2] it is only assumed that $M_\nu \geq 0$. As a consequence, in our case the 95% confidence regions for M_ν are shifted, by definition, towards larger masses. The same care should be applied when comparing to similar constraints reported in the literature.

The inclusion of BAO measurements results in much tighter neutrino mass bounds than those obtained with CMB data only. The combination of the cosmological data then starts to be sensitive to the region of m_{light} in which the NH and the IH scenarios correspond to different neutrino mass spectra. This can be clearly noticed from the bimodal distributions in both M_ν and in the neutrino mass fractions depicted in Fig. 1, where two




FIG. 1. One-dimensional probability posterior distribution of a selection of parameters analyzed in this work, for the combinations of current CMB and BAO datasets reported in the top legend. In the top panels, we report the posterior distributions for the sum of the neutrino masses $M_\nu = \Sigma_i m_i$, where the index $i = 1, 2, 3$ runs over the three mass eigenstates m_i ; the mass carried by the lightest eigenstate m_{light} and the Hubble constant H_0 . In the bottom panels, we report the posterior distributions for the neutrino mass fractions $f_{\nu,i} = m_i/M_\nu$. The solid (dashed) lines are for CMB alone (CMB plus BAO) measurements. The vertical dashed lines in the bottom panels refer to the expected value of $f_{\nu,i} = 1/3$ in the case of a fully degenerate mass spectrum. All the posterior shown in this figure also take into account information from oscillation measurements.

distinct peaks appear, each one being associated to the most probable value of the parameter *for a given choice of the hierarchy*. In fact, focusing on M_ν , we find that the most probable value is $M_\nu = 0.059 \text{ eV} = M_{\nu,\text{min}}^{\text{NH}}$, but a second peak is also clearly visible in $M_\nu = 0.098 \text{ eV} = M_{\nu,\text{min}}^{\text{IH}}$. These peaks are associated to the (single) peak at $m_{\text{light}} = 0$ in the posterior for the mass of the lightest neutrino, that gets mapped to two distinct values of M_ν , depending on the hierarchy.

By focusing on the one-dimensional distribution of the *hyperparameter* h_{type} , one might be able to assess whether current cosmological data favour one of the two hierarchical scenarios and to what extent. After the inspection of Fig. 1, upper panel, we can conclude that while there is still not compelling evidence for the cosmological data to prefer one of the two scenarios, the combination of CMB and BAO slightly favors the NH scheme (4:3 odds in favor of NH, without using small-

scale polarization, or 3:2 if we use it).

Finally, we note that the mean values and errors of the standard Λ CDM parameters $\Omega_c h^2$, H_0 and Ω_m shown in Tab. I are indeed very close to those quoted in Ref. [2] for the corresponding data sets, and derived without taking into account the uncertainty on the hierarchy⁷. This is yet another reflection of the fact that in the most part of the high-probability region of the parameter space, the mass spectrum is effectively degenerate.

To conclude this section, current cosmological data are only mildly sensitive to the neutrino mass ordering, with a preference of 4:3 in favour of NH from the combination

⁷ The cosmological parameters constraints for the Λ CDM model and many extensions, from several combinations of Planck 2015 and external data, can be downloaded from the Planck Legacy Archive <http://www.cosmos.esa.int/web/planck/pla>.

of Planck CMB data and BAO measurements.

IV. FORECASTS FOR FUTURE CMB AND BAO SURVEYS

In this section, we present forecasted constraints from future cosmological surveys on the neutrino mass parameters and discuss whether the improved sensitivity of the next-generation cosmological observatories will help unraveling the dilemma about the neutrino mass ordering.

We present the results for the forecasted COrE-like [32] CMB mission and a DESI-like survey [37] in Tab. II and Fig. 2. The results are quoted following the same scheme adopted in Tab. I and detailed at the beginning of Sec. III. We have considered three possible fiducial scenarios: two NH schemes, one with $M_\nu = 0.06$ eV and the other one with $M_\nu = 0.1$ eV, and one IH scenario, also with $M_\nu = 0.1$ eV. Notice that even for a future CMB mission as COrE it will be very difficult to extract with high statistical significance the neutrino mass hierarchy in any of the three fiducial scenarios explored here, even in the case of $m_{\text{light}} = 0$ ($M_\nu = 0.06$ eV). Nevertheless, by a comparison between the results in Tabs. I and those in Tab. II one can learn that the expected sensitivity of COrE alone on the neutrino mass measurements is slightly better than current combined CMB and galaxy clustering searches. This accuracy could be reinforced, for instance, by improved measurements from low-redshift experiments, such as adding a prior on the Hubble constant which simulates a $\sim 1\%$ -measurement of H_0 . However, the impact of such a constraint will be almost negligible, as the COrE mission alone reaches already that precision in H_0 . We therefore focus on additional information coming from galaxy surveys and add future forecasted measurements of the Hubble parameters and of the angular diameter distance from the DESI survey [37].

Adding BAO measurements improves considerably the results for the fiducial model with $M_\nu = 0.06$ eV; in this case, we find a 9 : 1 preference of NH versus IH. The great improvement due to the addition of DESI BAO data when $M_\nu = 0.06$ eV can be clearly visualized from the first two panels of Fig. 2: notice that the second peak at ~ 0.1 eV in the M_ν posterior is significantly reduced after the inclusion of BAO data.

On the contrary, for the $M_\nu = 0.1$ eV case, the situation is dramatically different. In fact, even if the addition of BAO measurements helps at pinpointing the value of sum of neutrino masses (as it can be seen by comparing, in Fig. 2, the width of the dashed black and blue curves with that of their solid counterparts), nevertheless the data still remain completely uninformative for what concerns the mass splitting, as it can be inferred by looking at the numbers reported in the last row of Tab. II, second and third columns. This points to the fact that, as already explained in Sec. II in reference to current experiments, the capability of future CMB and BAO observa-

tions to discriminate the neutrino mass hierarchy mainly relies on volume effects, i.e., on the possibility of excluding $M_\nu \geq 0.1$ eV with a high statistical significance; this is the case for the fiducial model with $M_\nu \geq 0.06$ eV. When instead $M_\nu = 0.1$ eV (or larger), as in the other two fiducial models considered here, the two mass orderings should be disentangled through the effect of the individual neutrino masses on the evolution of cosmological perturbations. Our findings clearly indicate that this is beyond the reach of next-generation CMB and BAO experiments, even in the most optimistic case (for $M_\nu > 0.1$ eV, the differences between the two hierarchies are even smaller). A possible improvement could come from highly accurate measurements of the matter power spectrum [11]; see also Ref. [40] for an appraisal of future 21 cm facilities. Alternatively, one should combine results coming from cosmological analysis with constraints obtained in laboratory searches, as we shall see in the following section.

V. IMPLICATIONS FOR $0\nu 2\beta$

In the following, we shall discuss the implications of our analyses for current and future neutrinoless double beta decay searches, as well as discuss the possibility to gain more information by the combination of cosmological observations with $0\nu 2\beta$ experiments. The non-observation of neutrinoless double β decay processes provides at present bounds on the so-called *effective Majorana mass* of the electron neutrino

$$m_{\beta\beta} = \frac{m_e}{\mathcal{M}\sqrt{G_{0\nu}T_{1/2}^{0\nu}}}, \quad (6)$$

where $T_{1/2}^{0\nu}$ is the neutrinoless double β decay half-life, m_e is the electron mass, $G_{0\nu}$ is a phase-space factor and \mathcal{M} is the nuclear matrix element (NME), a crucial quantity whose uncertainties affect significantly the interpretation of current and future searches for $0\nu 2\beta$ decay events.

The effective Majorana mass is related to the neutrino mass eigenvalues as follows:

$$m_{\beta\beta} \equiv \left| \sum_k U_{ek}^2 m_k \right| = \left| \sum_k e^{i\phi_k} V_{ek}^2 m_k \right|, \quad (7)$$

where we have written the mixing matrix U of Eq. (1) as the product of a matrix V , that contains the mixing angles and the Dirac phase, and a diagonal matrix that contains the *Majorana phases* ϕ_k , see e.g. Sec. 14 of Ref. [1]. Since one of the phases can always be rotated away, we can assume that $\phi_1 = 0$. While the elements of V are the same that enter the oscillation probabilities, the Majorana phases play no role in neutrino oscillation processes. However, they enter in the determination of the $0\nu 2\beta$ half-life, as it is clear from Eqs. (6)-(7), and are thus crucial for $0\nu 2\beta$ experiments. In fact, if the hierarchy is normal, the phases could also arrange to produce a

	COrE: $M_\nu = 0.1$ eV, NH (+ DESI)	COrE: $M_\nu = 0.1$ eV, IH (+ DESI)	COrE: $M_\nu = 0.06$ eV (+ DESI)
M_ν [eV]	$[0.058 - 0.188]$ $(0.112^{+0.037}_{-0.040})$	$[0.058 - 0.186]$ $(0.113^{+0.038}_{-0.042})$	$[0.058 - 0.155]$ $([0.058 - 0.109])$
m_{light} [eV]	< 0.0529 (< 0.0362)	< 0.0523 (< 0.0366)	< 0.0405 (< 0.0225)
m_1 [eV]	< 0.067 $([0.002 - 0.061])$	< 0.068 $([0.002 - 0.061])$	< 0.0571 (< 0.0491)
m_2 [eV]	$[0.009 - 0.0664]$ $([0.009 - 0.0555])$	$[0.009 - 0.0659]$ $([0.009 - 0.0562])$	$[0.009 - 0.0577]$ $([0.009 - 0.0499])$
m_3 [eV]	< 0.070 (< 0.061)	< 0.070 (< 0.062)	< 0.063 (< 0.055)
$\Omega_c h^2$	$0.1209^{+0.0011}_{-0.0010}$ (0.12072 ± 0.00058)	$0.1209^{+0.0010}_{-0.00098}$ $(0.12071^{+0.00058}_{-0.00057})$	0.12117 ± 0.00087 $(0.12079^{+0.00054}_{-0.00052})$
H_0 [km s $^{-1}$ Mpc $^{-1}$]	$66.27^{+0.95}_{-0.99}$ $(66.46^{+0.52}_{-0.48})$	$66.30^{+0.93}_{-0.97}$ $(66.46^{+0.53}_{-0.50})$	$[65.59 - 67.08]$ $(66.72^{+0.35}_{-0.44})$
Ω_m	$0.329^{+0.014}_{-0.013}$ $(0.3262^{+0.0066}_{-0.0070})$	$0.328^{+0.014}_{-0.013}$ (0.3262 ± 0.0070)	$[0.318 - 0.339]$ $(0.3230^{+0.0061}_{-0.0049})$
h_{type} odds	1 : 1 (1 : 1)	1 : 1 (1 : 1)	3 : 2 (9 : 1)

TABLE II. As Tab. I but for future measurements from the COrE CMB mission and for the DESI galaxy survey, and three different fiducial models: NH with $M_\nu = 0.1$ eV, IH with $M_\nu = 0.1$ eV and NH with $M_\nu = 0.06$ eV (second, third and fourth columns).

vanishing Majorana mass, and thus no observable $0\nu 2\beta$ signal, even for non vanishing values of the individual masses.

In the previous sections, we have combined cosmological observations (that mainly constrain M_ν) and oscillation measurements (that probe the mass differences) to derive limits on the individual masses, taking into account our ignorance of the mass hierarchy. Since oscillation measurements also constrain the elements of the mixing matrix, we can exploit the same strategy to derive constraints on the Majorana mass [related to the other parameters of the MCMC analysis by Eq. (7)], provided that we also take into account our ignorance of the true values of the Majorana phases.

Following Ref. [41], we thus consider the two Majorana phases as extra parameters in the MCMC analysis, ϕ_2 and ϕ_3 , with flat priors in the range $[0, 2\pi]$, as currently the values of these phases are totally unknown. We also do not consider here the possibility of future independent measurements of the phases. We then extend the analysis discussed in the previous sections for Planck TT,TE,EE+lowP+BAO, as an example of current data, and for COrE+DESI, as an example of future data, extracting the posterior distribution for the Majorana mass.

In case of future data, we consider two fiducial models with NH mass ordering and either $M_\nu = 0.06$ eV or $M_\nu = 0.1$ eV.

The 95% CL limit we find from current cosmological data, after marginalization over the *hyperparameter* h_{type} , is $m_{\beta\beta} < 0.056$ eV. We illustrate in Fig. 3 the 68% and 95% CL probability contours in the $M_\nu - m_{\beta\beta}$ plane. We also depict together the tightest bounds on neutrinoless double beta decay searches, coming from the KamLAND-Zen experiment, with 90% CL limits on $m_{\beta\beta} < 61 - 165$ meV, the precise value depending crucially on the NMEs assumption [42]. A visual inspection of the two-dimensional posterior makes evident that the posterior is bimodal, as it can be seen in Fig. 4. These results show that there exist two equally allowed possible regions for $m_{\beta\beta}$, due to the existing neutrino mass hierarchy ambiguity. Interestingly, current sensitivities to $m_{\beta\beta}$ start to reach the allowed region by cosmological and oscillation measurements, and, consequently, if nature has chosen the inverted hierarchy, a positive signal from neutrinoless double beta decay searches could be imminent (providing of course neutrinos possess a Majorana character and barring highly exotic physical scenarios). At present, both cosmological and laboratory tests of $m_{\beta\beta}$




FIG. 2. One-dimensional probability posterior distribution of a selection of parameters analyzed in this work, for the combination of datasets reported in the figure. In the top panels, we report the posterior distributions for the sum of the neutrino masses $M_\nu = \sum_i m_i$, where the index $i = 1, 2, 3$ runs over the three mass eigenstates m_i ; the mass carried by the lightest eigenstate m_{light} and the Hubble constant H_0 . In the bottom panels, we report the posterior distributions for the neutrino mass fractions $f_{\nu,i} = m_i/M_\nu$. The solid (dashed) lines refer to COre (COre plus DESI) forecasted MCMC results. The vertical dashed lines in the bottom panels refer to the expected value of $f_{\nu,i} = 1/3$ in the case of a fully degenerate mass spectrum. All the posterior shown in this figure also take into account information from oscillation measurements.

provide very similar constraints on the Majorana mass (assuming the most favorable values for the NMEs).

We would like to emphasize that the two equally-allowed regions in the $M_\nu/m_{\beta\beta}$ plane are not drawn separately from each other by assuming *a priori* one of the two hierarchies before performing the analysis, as usually done in literature. On the contrary, the appearance of the two regions is a direct consequence of the hierarchical model built via the *hyperparameter* h_{type} .

We have also performed a forecast to compute the expected sensitivities to $m_{\beta\beta}$ from future cosmological data. Combining COre and DESI, for a NH scenario, we obtain the 95% CL bounds of $m_{\beta\beta} < 0.034$ eV and $0.005 \text{ eV} < m_{\beta\beta} < 0.053$ eV, assuming that $M_\nu = 0.06$ eV and $M_\nu = 0.1$ eV, respectively. Notice that for the $M_\nu = 0.1$ eV case the expected limit on $m_{\beta\beta}$ is very close to the current one, as for this particular scenario future cosmological measurements will most likely be unable to determine the neutrino mass ordering. Figure 5 depicts the two-dimensional contours in the $M_\nu - m_{\beta\beta}$ plane for the two possible fiducial models above mentioned, together with the expected upper bounds (assuming $m_{\beta\beta} = 0$) from a future, nEXO-like [43] neutrinoless double beta decay experiment. In the case of

$M_\nu = 0.1$ eV the prospects of observing a positive signal from a future $0\nu 2\beta$ decay are very good (provided neutrinos are Majorana and the mass mechanism is responsible for $0\nu 2\beta$ decay), despite the fact that the hierarchy can not be determined via cosmological measurements. In this situation the hierarchy could be extracted by neutrinoless double beta decay itself, since a positive signal characterized by $m_{\beta\beta} \simeq 0.05$ eV would suggest an IH scenario. Alternatively, a positive signal characterized by $m_{\beta\beta} \simeq 0.02$ eV plus the expected sensitivity of $\sigma(m_{\beta\beta}) \sim 0.01$ eV would point to a NH scenario and discard the IH scenario with high statistical significance.

If, on the other hand, M_ν is closer to the minimal value allowed in the NH scenario, the sensitivity of future neutrinoless double beta decay searches may not be enough to detect the putative signal from Majorana neutrinos, due to the possible disruptive interference played by oscillation parameters in the definition of the Majorana mass. Nevertheless, in that case, future cosmological data can single out the neutrino mass ordering with high significance.

VI. CONCLUSIONS

We have presented constraints on cosmological parameters in the context of a Λ CDM + M_ν scenario, with M_ν representing the sum of neutrino masses, assuming three massive non-degenerate eigenstates and properly taking into account the neutrino mass ordering. Indeed, the novelty of the study presented here relies on our treatment of the neutrino mass ordering, currently totally unknown. We implement the neutrino hierarchy ambiguity by means of a *hyperparameter* h_{type} to be marginalized over. This approach allows us to (i) model the exact mass splittings without making use of approximations, and including the information from oscillation measurements; (ii) to quantitatively assess the preference for one of the two hierarchies in a straightforward fashion, without the need for computing the Bayesian evidence for performing model comparison, and (iii) to account for the incomplete knowledge of the neutrino hierarchy that could potentially affect the neutrino mass bounds. We have employed current cosmological data coming from the Planck satellite measurements of the CMB anisotropies and a compilation of BAO measurements at different redshifts. We have also performed forecasts for future cosmological missions, such as the proposed CMB satellite mission COrE and the future galaxy survey DESI.

Focusing on current cosmological measurements, we have shown that CMB temperature and polarization data alone are not sensitive enough to discriminate between the two hierarchies. When BAO information is included, present cosmological probes start to be weakly sensitive to the mass ordering (3 : 2 or 4 : 3 odds in favor of NH, with or without CMB small-scale polarization, respectively), although compelling evidence for one of the two is still lacking. Marginalizing over the hierarchy parameter slightly worsens the neutrino mass limits, albeit galaxy clustering data in the form of BAO measurements lead to results very similar to those obtained in the absence of the *hyperparameter* h_{type} .

Concerning future experiments, their combination turns out to be really powerful, as it will lead to a 9 : 1 preference for the normal hierarchy scenario versus the inverted hierarchy one, assuming a fiducial cosmology with a sum of neutrino masses $M_\nu = 0.06$ eV. However, for larger masses $M_\nu = 0.1$ eV distinguishing the hierarchy via cosmological measurements alone turns out to be an extremely difficult task. Adding other possible future improvements, as, for instance, a 1% prior in the value of the Hubble constant, will not change significantly the results, since the COrE mission is expected to provide a smaller uncertainty on H_0 . Additional constraining power might come from more precise measurements of the shape of the matter power spectrum, provided that systematics and uncertainties related to the exact modelling of the perturbation behaviour in the non-linear regime (where we expect neutrinos to leave their most peculiar signature on the matter power spectrum) are kept under control.




FIG. 3. Two dimensional 68% and 95% probability contours in the $M_\nu - m_{\beta\beta}$ plane, for current cosmological (Planck TT, TE, EE + BAO) and neutrino oscillation data. The contours are calculated by taking into account the uncertainty on the mass ordering, without fixing a priori the mass hierarchy. Two distinct contour areas, corresponding to the two mass orderings, can be identified (even if they are not completely separated from each other), see text for details. The orange horizontal bands correspond to the 90% upper bounds on $m_{\beta\beta}$ obtained from KamLAND-Zen [42], for different assumptions for the values of the nuclear matrix elements that enter into the calculation of $m_{\beta\beta}$.

We have also studied the implications and the complementarity with neutrinoless double beta decay searches. Current limits from the KamLAND-Zen experiment are competitive and consistent with the tightest cosmological limit we find here on the *effective Majorana mass*, $m_{\beta\beta} < 0.056$ eV. These results imply that, if nature has chosen the inverted hierarchy scheme and the Majorana neutrino character (versus the normal hierarchy scenario and the Dirac nature), a positive signal from neutrino neutrinoless double beta decay searches, as well as a cosmological detection of the neutrino mass, could be imminent. Future prospects for neutrinoless double beta decay experiment are promising for a total neutrino mass $M_\nu = 0.1$ eV, regardless of the neutrino mass hierarchy. However, if the lightest neutrino mass eigenstate turns out to be zero, and the hierarchy normal, the detection of this putative signal cannot be guaranteed, even for an ultimate, highly sensitive neutrinoless double beta decay experiment. The good news is that if this is the case realized in nature, cosmology will be able to tell us about the neutrino mass hierarchy with compelling statistical significance.




FIG. 4. Density plot of the two-dimensional posterior in the $M_\nu - m_{\beta\beta}$ plane, for current cosmological (Planck TT, TE, EE + BAO) and neutrino oscillation data. Darker colors correspond to higher probability regions. The plot shows that the posterior is bimodal.

ACKNOWLEDGMENTS

Based on observations obtained with Planck (<http://www.esa.int/Planck>), an ESA science mission with instruments and contributions directly funded by ESA Member States, NASA, and Canada. We also acknowledge use of the Planck Legacy Archive. M.G. and K.F. are supported by the Vetenskapsrådet (Swedish Research Council, Contract No. 638-2013-8993). M.L. acknowledges support from ASI through ASI/INAF Agreement 2014-024-R.1 for the Planck LFI Activity of Phase E2. O.M. is supported by PROMETEO II/2014/050, by the Spanish Grant FPA2014-57816-P of the MINECO, by the MINECO Grant SEV-2014-0398 and by the European Unions Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreements 690575 and 674896. K.F. acknowledges support by the DoE grant de-sc0007859.

-
- [1] C. Patrignani *et al.* [Particle Data Group], *Chin. Phys. C*, **40**, 100001 (2016).
 - [2] P. A. R. Ade *et al.* [Planck Collaboration], *Astron. Astrophys.* **594**, A13 (2016) doi:10.1051/0004-6361/201525830 [arXiv:1502.01589 [astro-ph.CO]].
 - [3] N. Palanque-Delabrouille *et al.*, *JCAP* **1511**, no. 11, 011 (2015) doi:10.1088/1475-7516/2015/11/011 [arXiv:1506.05976 [astro-ph.CO]].
 - [4] E. Di Valentino, E. Giusarma, O. Mena, A. Melchiorri and J. Silk, *Phys. Rev. D* **93**, no. 8, 083527 (2016) doi:10.1103/PhysRevD.93.083527 [arXiv:1511.00975 [astro-ph.CO]].
 - [5] A. J. Cuesta, V. Niro and L. Verde, *Phys. Dark Univ.* **13**, 77 (2016) doi:10.1016/j.dark.2016.04.005 [arXiv:1511.05983 [astro-ph.CO]].
 - [6] E. Giusarma, M. Gerbino, O. Mena, S. Vagnozzi, S. Ho and K. Freese, arXiv:1605.04320 [astro-ph.CO].
 - [7] S. Hannestad and T. Schwetz, arXiv:1606.04691 [astro-ph.CO].
 - [8] J. Lesgourgues, S. Pastor and L. Perotto, *Phys. Rev. D* **70**, 045016 (2004) doi:10.1103/PhysRevD.70.045016 [hep-ph/0403296].
 - [9] A. Slosar, *Phys. Rev. D* **73**, 123501 (2006) doi:10.1103/PhysRevD.73.123501 [astro-ph/0602133].
 - [10] J. Hamann, S. Hannestad and Y. Y. Y. Wong, *JCAP* **1211**, 052 (2012) doi:10.1088/1475-7516/2012/11/052 [arXiv:1209.1043 [astro-ph.CO]].
 - [11] R. Jimenez, T. Kitching, C. Pena-Garay and L. Verde, *JCAP* **1005**, 035 (2010) doi:10.1088/1475-7516/2010/05/035 [arXiv:1003.5918 [astro-ph.CO]].
 - [12] L. Xu and Q. G. Huang, arXiv:1611.05178 [astro-ph.CO].
 - [13] F. T. Avignone, III, S. R. Elliott and J. Engel, *Rev. Mod. Phys.* **80**, 481 (2008) doi:10.1103/RevModPhys.80.481 [arXiv:0708.1033 [nucl-ex]].
 - [14] O. Cremonesi and M. Pavan, *Adv. High Energy Phys.* **2014**, 951432 (2014) doi:10.1155/2014/951432 [arXiv:1310.4692 [physics.ins-det]].
 - [15] J. J. Gomez-Cadenas, J. Martin-Albo, M. Mezzetto, F. Monrabal and M. Sorel, *Riv. Nuovo Cim.* **35**, 29 (2012) doi:10.1393/ncr/i2012-10074-9 [arXiv:1109.5515 [hep-ex]].
 - [16] S. Dell’Oro, S. Marocco, M. Viel and F. Visani, *Adv. High Energy Phys.* **2016**, 2162659 (2016) doi:10.1155/2016/2162659 [arXiv:1601.07512 [hep-ph]].
 - [17] J. Schechter and J. W. F. Valle, *Phys. Rev. D* **25**, 2951 (1982). doi:10.1103/PhysRevD.25.2951
 - [18] A. Gelman, J. B. Carlin, H. S. Stern, D. B. Dunson, A. Vehtari and D. B. Rubin, (2013), *Bayesian Data Analysis (Third edition)*, Chapman & Hall/CRC Texts in Statistical Science.
 - [19] [NOvA Collaboration], http://neutrino2016.iopconfs.org/IOP/media/uploaded/EVIOP/event_948/11.30_-_has_to_be_PDF.pdf
 - [20] <http://www.nu-fit.org> J. Bergstrom, M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, *JHEP* **1509**, 200 (2015) doi:10.1007/JHEP09(2015)200 [arXiv:1507.04366 [hep-ph]]; M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, *JHEP* **1411**, 052 (2014) doi:10.1007/JHEP11(2014)052 [arXiv:1409.5439 [hep-ph]].
 - [21] D. V. Forero, M. Tortola and J. W. F. Valle, *Phys. Rev. D* **90**, no. 9, 093006 (2014) doi:10.1103/PhysRevD.90.093006 [arXiv:1405.7540 [hep-ph]].




FIG. 5. Two dimensional 68% and 95% probability contours in the $M_\nu - m_{\beta\beta}$ plane, for future cosmological (CORe+DESI) and neutrino oscillation data, considering the NH scenario as the nature's choice and $M_\nu = 0.06$ eV (left panel) or $M_\nu = 0.1$ eV (right panel). The contours are calculated by taking into account the uncertainty on the mass ordering, without fixing a priori the mass hierarchy. The two regions that can be inferred in both figures (even if not completely isolated) correspond to the two mass orderings. The horizontal bands correspond to the 90% upper bounds on $m_{\beta\beta}$ obtained from KamLAND-Zen [42] (orange) and to those expected from a future nEXO-like experiment (green) [43] (assuming a vanishing Majorana mass), for different assumptions for the values of the nuclear matrix elements that enter into the calculation of $m_{\beta\beta}$.

- [22] A. Lewis and S. Bridle, Phys. Rev. D **66**, 103511 (2002) doi:10.1103/PhysRevD.66.103511 [astro-ph/0205436].
- [23] A. Lewis, Phys. Rev. D **87**, no. 10, 103529 (2013) doi:10.1103/PhysRevD.87.103529 [arXiv:1304.4473 [astro-ph.CO]].
- [24] S. Brooks and A. Gelman, J. Comp. Graph. Stat. **7**, 434-455 (1998).
- [25] R. Adam *et al.* [Planck Collaboration], doi:10.1051/0004-6361/201527101 arXiv:1502.01582 [astro-ph.CO].
- [26] N. Aghanim *et al.* [Planck Collaboration], Astron. Astrophys. doi:10.1051/0004-6361/201526926 [arXiv:1507.02704 [astro-ph.CO]].
- [27] D. J. Eisenstein and W. Hu, Astrophys. J. **496**, 605 (1998) doi:10.1086/305424 [astro-ph/9709112].
- [28] F. Beutler *et al.*, Mon. Not. Roy. Astron. Soc. **416**, 3017 (2011) doi:10.1111/j.1365-2966.2011.19250.x [arXiv:1106.3366 [astro-ph.CO]].
- [29] L. Anderson *et al.* [BOSS Collaboration], Mon. Not. Roy. Astron. Soc. **441**, no. 1, 24 (2014) doi:10.1093/mnras/stu523 [arXiv:1312.4877 [astro-ph.CO]].
- [30] E. Giusarma, R. De Putter and O. Mena, Phys. Rev. D **87**, no. 4, 043515 (2013) doi:10.1103/PhysRevD.87.043515 [arXiv:1211.2154 [astro-ph.CO]].
- [31] N. Aghanim *et al.* [Planck Collaboration], Astron. Astrophys. in press, [arXiv:1605.02985 [astro-ph.CO]].
- [32] F. R. Bouchet *et al.* [CORe Collaboration], arXiv:1102.2181 [astro-ph.CO].
- [33] J. R. Bond, A. H. Jaffe and L. E. Knox, Astrophys. J. **533**, 19 (2000) doi:10.1086/308625 [astro-ph/9808264].
- [34] J. R. Bond, G. Efstathiou and M. Tegmark, Mon. Not. Roy. Astron. Soc. **291**, L33 (1997) doi:10.1093/mnras/291.1.L33 [astro-ph/9702100].
- [35] J. Errard, S. M. Feeney, H. V. Peiris and A. H. Jaffe, JCAP **1603**, no. 03, 052 (2016) doi:10.1088/1475-7516/2016/03/052 [arXiv:1509.06770 [astro-ph.CO]].
- [36] M. Escudero, H. Ramrez, L. Boubekur, E. Giusarma and O. Mena, JCAP **1602**, no. 02, 020 (2016) doi:10.1088/1475-7516/2016/02/020 [arXiv:1509.05419 [astro-ph.CO]].
- [37] M. Levi *et al.* [DESI Collaboration], arXiv:1308.0847 [astro-ph.CO].
- [38] A. Font-Ribera, P. McDonald, N. Mostek, B. A. Reid, H. J. Seo and A. Slosar, JCAP **1405**, 023 (2014) doi:10.1088/1475-7516/2014/05/023 [arXiv:1308.4164 [astro-ph.CO]].
- [39] H. J. Seo and D. J. Eisenstein, Astrophys. J. **665**, 14 (2007) doi:10.1086/519549 [astro-ph/0701079].
- [40] J. R. Pritchard and E. Pierpaoli, Phys. Rev. D **78**, 065009 (2008) doi:10.1103/PhysRevD.78.065009 [arXiv:0805.1920 [astro-ph]].
- [41] M. Gerbino, M. Lattanzi and A. Melchiorri, Phys. Rev. D **93**, no. 3, 033001 (2016) doi:10.1103/PhysRevD.93.033001 [arXiv:1507.08614 [hep-ph]].

- [42] A. Gando *et al.* [KamLAND-Zen Collaboration], Phys. Rev. Lett. **117**, no. 8, 082503 (2016) Addendum: [Phys. Rev. Lett. **117**, no. 10, 109903 (2016)] doi:10.1103/PhysRevLett.117.109903, 10.1103/Phys-RevLett.117.082503 [arXiv:1605.02889 [hep-ex]].
- [43] <https://www-project.slac.stanford.edu/exo/>